

Product	Active ingredient	Rate / A	REI Hrs.	PHI Days	Efficacy (USDA)	Max. per crop / season	App. Interval
Actara 25WDG	Thiamethoxam	2.0-5.5 oz/A	12	35	+++	16.5 oz./A (0.258 lb. a.i./A)	10d
Asana XL 0.66EC	Esfenvalerate	4.8-14.5 fl oz/A	12	21	++	101 fl oz/A (0.525 lb AI/A).	NA
Baythroid XL 1EC	Beta-Cyfluthrin	1.4-2.8 fl oz/A	12	7	++	2.8 fl oz/A (0.022 lb AI/A).	14d
Bifenture EC	Bifenthrin	5.2-12.8 fl oz/A	12	14	++++	32 fl ozs (0.50 lbs ai)	30d
Bifenture 10DF	Bifenthrin	12.8-32.0 oz/A	12	14	++++	80 ozs (0.50 lbs ai)	30d
Brigade WSB	Bifenthrin	12.8-32.0 oz/A	12	14	++++	80 ozs (0.50 lbs ai)	30d
Danitol 2.4EC	Fenpropathrin	10.66-21.33 fl oz/A	24	14	+++	42.56 fl ozs (0.80 lbs ai)	10d
Endigo ZC	Thiamethoxam / Lambda-cyhalothrin	5-6 fl fl oz/A	24	35	++++	19 fl oz./A (0.172 lb ai) NY	10d
Lannate 2.4LV*	Methomyl	2.25 pt/A	72	14	++++	240 ozs (0.50 lbs ai)	7d
Lannate 90SP*	Methomyl	8-16 oz/A	72	14	++++	5.0 lbs	7d
Leverage 360	Beta-Cyfluthrin / Imidacloprid	2.4-2.8 fl oz/A	12	7	+++	2.8 fl oz/A	14d
Surround 95WP	Kaolin	25-50 lb/A	4	0	+	NA	0d
Thionex 50WP	Endosulfan	Max. 5 lb/A	20 days	21	++++	6.0 lbs	NA
Thionex EC	Endosulfan	1.33-2.67 qts./A	7 days	21	++++	2-2/3 qts (2.0 lbs ai)	NA
Voliam Xpress EC	Chlorantraniliprole / Lambda-cyhalothrin	6-12 fl oz/A	24	21	+++	31.0 fl oz/A	10d
Vydate 2L*	Oxamyl	4-8 pt/A	48	14	++	281 fl oz/A (128 oz AI/A).	7d
Warrior 1CS	Lambda-cyhalothrin	2.56-5.12 fl oz/A	24	21	++	20.48 fl. oz. (0.28 lb. a.i.)*	5d
Warrior II 2.08CS	Lambda-cyhalothrin	1.28-2.56 fl oz/A	24	21	++	10.24 fl. oz. (0.28 lb. a.i.)*	5d

\* Although these materials have excellent topical ratings in lab bioassay studies, field efficacy studies have shown economic fruit injury from BMSB feeding, suggesting low residual levels.

\*\* Post bloom applications

(+) low to (++++) high efficacy