

June 2017 - Labeled Insecticides for Control of Spotted Wing Drosophila in New York Berry Crops – Quick Guide

Compiled by Greg Loeb, Laura McDermott, Peter Jentsch, Tess Grasswitz, & Juliet Carroll, Cornell University. Updated regularly.

BLUEBERRIES										
PRODUCT	AI ¹	IRAC group ²	EPA#	Rate/A	REI ³	DTH ⁴	Max. Prod/A/yr (ai)	Total applic's	Spray Interval	Probable efficacy
^@Entrust Naturalyte (2ee) ^a	spinosad	5	62719-282	1.25-2 oz	4 hr	3 d	9 oz (0.45 lb)	3 per crop	> 6 d	Good to Excellent [#]
^@Entrust SC ^a	spinosad	5	62719-621	4-6 fl oz	4 hr	1 d	29 fl oz (0.45 lb)	3 per crop	> 6 d	Good to Excellent [#]
@Delegate WG (2ee)	spinetoram	5	62719-541	3-6 oz	4 hr	3 d	19.5 oz (0.305 lb)	6	> 6 d	Excellent [#]
*Exirel	cyazypyr	28	352-859	13.5-20.5 fl oz	12 hr	3 d	61.5 fl oz (0.4 lb)	3	> 5 d	Excellent
*Brigade WSB (2ee)	bifenthrin	3A	279-3108	5.3-16 oz	12 hr	1 d	5 lb (0.5 lb)	-	> 7 d	Excellent
*Danitol 2.4EC	fenpropathrin	3A	59639-35	16 fl oz	24 hr	3 d	32 fl oz (0.6 lb)	2	-	Excellent
*Mustang Max Insecticide (2ee)	zeta-cypermethrin	3A	279-3249	4 fl oz	12 hr	1 d	24 fl oz (0.15 lb)	6	> 7 d	Excellent
*Mustang Maxx Insecticide (2ee)	zeta-cypermethrin	3A	279-3426	4 fl oz	12 hr	1 d	24 fl oz (0.15 lb)	6	> 7 d	Excellent
^Pyganic EC 1.4	pyrethrin	3A	1021-1771	1 pt-2 qts	12 hr	0 d	-	-	-	Fair to Poor
^Pyganic EC 5.0	pyrethrin	3A	1021-1772	4.5-18 fl oz	12 hr	0 d	-	-	-	Fair to Poor
*Triple Crown	bifenthrin, imidacloprid, zeta-cypermethrin	3A,4A	279-3440	6.4-10.3 fl oz	12 hr	3 d	31 fl oz (0.54 lb)	5	> 7 d	Good to Excellent
Assail 30SG	acetamiprid	4A	8033-36-70506	4.5-5.3 oz	12 hr	1 d	26.7 oz (0.5 lb)	5	> 7 d	Good [#]
*Lannate SP	methomyl	1A	352-342	0.5-1 lb	48 hr	3 d	4 lb (3.6 lb)	4	> 5-7 d	Excellent

*Refer to label for details and additional restrictions.

#Adding sugar (sucrose) at 2 lb/100 gal water as a feeding stimulant will increase efficacy.

^Approved for organic use in NY.

@After two consecutive applications must rotate to different mode of action.

¹ Active Ingredient.

² Mode of Action, based on IRAC group code (UN = unknown).

³ Re-entry Interval (hr = hours).

⁴ Days to Harvest (d = days).

June 2017 - Labeled Insecticides for Control of Spotted Wing Drosophila in New York Berry Crops – Quick Guide

Compiled by Greg Loeb, Laura McDermott, Peter Jentsch, Tess Grasswitz, & Juliet Carroll, Cornell University. Updated regularly.

BLUEBERRIES										
PRODUCT	AI ¹	IRAC group ²	EPA#	Rate/A	REI ³	DTH ⁴	Max. Prod/A/yr (ai)	Total applic's	Spray Interval	Probable efficacy
*Lannate VP	methomyl	1A	352-384	1.5-3 pts	48 hr	3 d	12 pts (3.6 lb)	4	> 5-7 d	Excellent
*Imidan 70W	phosmet	1B	10163-169	1.33 lb	24 hr	3 d	7.125 lb (5.0 lb)	5	-	Excellent
Malathion 5EC (2ee)	malathion	1B	19713-217	2 pts	12 hr	1 d	6 pts (3.75 lb)	3	> 5 d	Good
Malathion 5EC (2ee)	malathion	1B	66330-220	2 pts	12 hr	1 d	6 pts (3.75 lb)	3	> 5 d	Good
Malathion 8 Aquamul (2ee)	malathion	1B	34704-474	2.5 pts	12 hr	1 d	3.75 pts (3.75 lb)	1	> 5 d	Good
Malathion 57 (2ee)	malathion	1B	67760-40-53883	2 pts	12 hr	1 d	6 pts (3.75 lb)	3	> 5 d	Good
^AzaSol	azadirachtin	UN	81899-4	6 oz in 50 gal	4 hr	0 d	-	-	-	Fair to Poor
^#Grandevo	<i>Chromobacterium subtsugae</i> strain PRAA4-1 and spent fermentation media	UN	84059-27	2-3 lb	4 hr	0 d	-	-	≤ 7 d	Fair to Poor

^a In organic production, Entrust must be rotated with insecticides with different modes of action, consider using Grandevo or products containing the active ingredients azadirachtin or pyrethrin.

*Refer to label for details and additional restrictions.

#Adding sugar (sucrose) at 2 lb/100 gal water as a feeding stimulant will increase efficacy.

^Approved for organic use in NY.

@After two consecutive applications must rotate to different mode of action.

¹ Active Ingredient.

² Mode of Action, based on IRAC group code (UN = unknown).

³ Re-entry Interval (hr = hours).

⁴ Days to Harvest (d = days).

June 2017 - Labeled Insecticides for Control of Spotted Wing Drosophila in New York Berry Crops – Quick Guide

Compiled by Greg Loeb, Laura McDermott, Peter Jentsch, Tess Grasswitz, & Juliet Carroll, Cornell University. Updated regularly.

RASPBERRIES & BLACKBERRIES										
PRODUCT	AI ¹	IRAC group ²	EPA#	RATE/A	REI ³	DTH ⁴	Max. Prod/A/yr (ai)	Total applic's	Spray Interval	Probable efficacy
[^] @Entrust Naturalyte (2ee) ^a	spinosad	5	62719-282	1.25-2 oz	4 hr	1 d	9 oz (0.45 lb)	3 per crop	> 6 d	Good to Excellent [#]
[^] @Entrust SC (2ee) ^a	spinosad	5	62719-621	4-6 fl oz	4 hr	1 d	29 fl oz (0.45 lb)	3 per crop	> 6 d	Good to Excellent [#]
[@] Delegate WG (2ee)	spinetoram	5	62719-541	3-6 oz	4 hr	1 d	19.5 oz (0.305 lb)	6	> 4 d	Excellent [#]
*Brigade WSB (2ee)	bifenthrin	3A	279-3108	8-16 oz	12 hr	3 d	2 lb (0.2 lb)	1 post bloom	-	Excellent
*Brigade EC (2ee)	bifenthrin	3A	279-3313	3.2-6.4 fl oz	12 hr	3 d	12.8 fl oz (0.2 lb)	1 post bloom	-	Excellent
*Danitol 2.4EC	fenpropathrin	3A	59639-35	16 fl oz	24 hr	3 d	32 fl oz (0.6 lb)	2	-	Excellent
*Mustang Max Insecticide (2ee)	zeta-cypermethrin	3A	279-3249	4 fl oz	12 hr	1 d	24 fl oz (0.15 lb)	6	> 7 d	Excellent
*Mustang Maxx Insecticide (2ee)	zeta-cypermethrin	3A	279-3426	4 fl oz	12 hr	1 d	24 fl oz (0.15 lb)	6	> 7 d	Excellent
[^] Pyganic EC 1.4	pyrethrin	3A	1021-1771	1 pt-2 qts	12 hr	0 d	-	-	-	Fair to Poor
[^] Pyganic EC 5.0	pyrethrin	3A	1021-1772	4.5-18 fl oz	12 hr	0 d	-	-	-	Fair to Poor
*Triple Crown	bifenthrin, imidacloprid, zeta-cypermethrin	3A, 4A	279-3440	6.4-10.3 fl oz	12 hr	3 d	10.3 fl oz (0.181 lb)	1 post bloom	> 7 d	Good to Excellent
Assail 30SG	acetamiprid	4A	8033-36-70506	4.5-5.3 oz	12 hr	1 d	26.7 oz (0.5 lb)	5	> 7 d	Good [#]
Malathion 5EC (2ee)	malathion	1B	19713-217	3 pts	12 hr	1 d	9 pts (6.0 lb)	3	> 7 d	Good

*Refer to label for details and additional restrictions.

[#]Adding sugar (sucrose) at 2 lb/100 gal water as a feeding stimulant will increase efficacy.

[^]Approved for organic use in NY.

[@]After two consecutive applications must rotate to different mode of action.

¹ Active Ingredient.

² Mode of Action, based on IRAC group code (UN = unknown).

³ Re-entry Interval (hr = hours).

⁴ Days to Harvest (d = days).

June 2017 - Labeled Insecticides for Control of Spotted Wing Drosophila in New York Berry Crops – Quick Guide

Compiled by Greg Loeb, Laura McDermott, Peter Jentsch, Tess Grasswitz, & Juliet Carroll, Cornell University. Updated regularly.

RASPBERRIES & BLACKBERRIES										
PRODUCT	AI ¹	IRAC group ²	EPA#	RATE/A	REI ³	DTH ⁴	Max. Prod/A/yr (ai)	Total applic's	Spray Interval	Probable efficacy
Malathion 5EC (2ee)	malathion	1B	66330-220	3 pts	12 hr	1 d	9 pts (6.0 lb)	3	> 7 d	Good
Malathion 8 Aquamul (2ee)	malathion	1B	34704-474	2 pts	12 hr	1 d	6 pts (6.0 lb)	3	> 7 d	Good
Malathion 57 (2ee)	malathion	1B	67760-40-53883	3 pts	12 hr	1 d	9 pts (6.0 lb)	3	> 7 d	Good
[^] AzaSol		UN	81899-4	6 oz in 50 gal	4 hr	0 d	-	-	-	Fair to Poor
Molt-X	azadirachtin	UN	68539-11	10 oz in 50 gal	4 hr	0 d	-	-	-	Fair to Poor
[^] #Grandevo	<i>Chromobacterium subtsugae</i> strain PRAA4-1 and spent fermentation media	UN	84059-27	2-3 lb	4 hr	0 d	-	-	≤ 7 d	Fair to Poor

^a In organic production, Entrust must be rotated with insecticides with different modes of action, consider using Grandevo or products containing the active ingredients azadirachtin or pyrethrin.

*Refer to label for details and additional restrictions.

#Adding sugar (sucrose) at 2 lb/100 gal water as a feeding stimulant will increase efficacy.

[^]Approved for organic use in NY.

[@]After two consecutive applications must rotate to different mode of action.

¹ Active Ingredient.

² Mode of Action, based on IRAC group code (UN = unknown).

³ Re-entry Interval (hr = hours).

⁴ Days to Harvest (d = days).

June 2017 - Labeled Insecticides for Control of Spotted Wing Drosophila in New York Berry Crops – Quick Guide

Compiled by Greg Loeb, Laura McDermott, Peter Jentsch, Tess Grasswitz, & Juliet Carroll, Cornell University. Updated regularly.

STRAWBERRIES										
PRODUCT	AI ¹	IRAC group ²	EPA#	RATE/A	REI ³	DTH ⁴	Max. Prod/A/yr (ai)	Total applic's	Spray Interval	Probable efficacy
^@Entrust Naturalyte (2ee) ^a	spinosad	5	62719-282	1.25-2 oz	4 hr	1 d	9 oz (0.45 lb)	5	> 5 d	Good to Excellent [#]
^@Entrust SC (2ee) ^a	spinosad	5	62719-621	4-6 fl oz	4 hr	1 d	29 fl oz (0.45 lb)	5	> 5 d	Good to Excellent [#]
@Radiant (2ee)	spinetoram	5	62719-545	6-10 fl oz	4 hr	1 d	39 fl oz (0.305 lb)	5	> 3 d	Excellent [#]
*Brigade WSB (2ee)	bifenthrin	3A	279-3108	8-16 oz	12 hr	0 d	5 lb (0.5 lb)	-	> 7 d	Excellent
*Danitol 2.4EC	fenpropathrin	3A	59639-35	16-21.3 fl oz	24 hr	2 d	42.7 fl oz (0.8 lb)	2	-	Excellent
^Pyganic EC 1.4	pyrethrin	3A	1021-1771	1 pt-2 qts	12 hr	0 d	-	-	-	Fair to Poor
^Pyganic EC 5.0	pyrethrin	3A	1021-1772	4.5-18 fl oz	12 hr	0 d	-	-	-	Fair to Poor
Assail 30SG	acetamiprid	4A	8033-36-70506	4.5-5.3 oz	12 hr	1 d	13.8 oz (0.26 lb)	2	> 7 d	Good [#]
Malathion 5EC (2ee)	malathion	1B	19713-217	3.2 pts	12 hr	3 d	12.8 pts (8 lb)	4	> 7 d	Good
Malathion 5EC (2ee)	malathion	1B	66330-220	3.2 pts	12 hr	3 d	12.8 pts (8 lb)	4	> 7 d	Good
Malathion 8 Aquamul (2ee)	malathion	1B	34704-474	2.0 pts	12 hr	3 d	8 pts (8 lb)	4	> 7 d	Good
Malathion 57 (2ee)	malathion	1B	67760-40-53883	3.2 pts	12 hr	3 d	12.8 pts (8 lb)	4	> 7 d	Good
^AzaSol	azadirachtin	UN	81899-4	6 oz in 50 gal	4 hr	0 d	-	-	-	Fair to Poor
^#Grandevo	<i>Chromobacterium subsugae</i> strain PRAA4-1 and spent fermentation media	UN	84059-27	2-3 lb	4 hr	0 d	-	-	≤ 7 d	Fair to Poor

^a In organic production, Entrust must be rotated with insecticides with different modes of action, consider using Grandevo or products containing the active ingredients azadirachtin or pyrethrin.

*Refer to label for details and additional restrictions.

[#]Adding sugar (sucrose) at 2 lb/100 gal water as a feeding stimulant will increase efficacy.

[^]Approved for organic use in NY.

[@]After two consecutive applications must rotate to different mode of action.

¹ Active Ingredient.

² Mode of Action, based on IRAC group code (UN = unknown).

³ Re-entry Interval (hr = hours).

⁴ Days to Harvest (d = days).